

# 5s correlation confinement resonances in Xe-endo-fullerenes

V. K. Dolmatov<sup>1</sup> and D. A. Keating

Department of Physics and Earth Science, University of North Alabama, Florence, AL 35632, U.S.A.


**Synopsis** Spectacular trends in the modification of the Xe 5s photoionization via interchannel coupling with confinement resonances emerging in the Xe 4d giant resonance upon photoionization of the Xe@C<sub>60</sub>, Xe@C<sub>240</sub> and Xe@C<sub>60</sub>@C<sub>240</sub> endo-fullerenes are theoretically unraveled and interpreted.

Resonances, termed *confinement resonances* (CR's), emerging in photoionization spectra of atoms  $A$  confined inside the empty spaces of endo-fullerenes -  $A@C_n$ ,  $A@C_n@C_m$ , etc - have attracted much attention of investigators in recent years [1] (and references therein). CR's occur in endo-fullerene photoionization spectra due to the interference of the photoelectron waves emerging directly from the confined atom  $A$ , and those scattered off the endo-fullerene confining cage(s). Their existence has recently been confirmed experimentally [2]. *Correlation confinement resonances* (CCR's) differ from ordinary CR's in that they occur in the photoionization of an *outer* subshell of the confined atom  $A$  due to the interference of transitions from the outer subshell with ordinary CR's emerging in *inner* shell photoionization transitions, via interchannel coupling [1, 3]. They were originally theoretically discovered in the Xe 5s photoionization of Xe@C<sub>60</sub> [3]. The CCR phenomenon thus introduces a novel class of resonances that can exist neither without a confinement nor electron correlation, thereby attracting much interest to their study.

Ordinary CR's in the Xe 4d giant photoionization resonance undergo dramatic changes along the path from Xe@C<sub>60</sub> to Xe@C<sub>240</sub> to Xe@C<sub>60</sub>@C<sub>240</sub> [4]. With the impetus of results of Ref. [4], we study how ordinary CR's in the Xe 4d photoionization affect the Xe 5s photoionization in various Xe-endo-fullerenes. It is the aim of this paper to report on spectacular CCR's emerging in the Xe 5s photoionization cross section due to interchannel coupling of the 5s  $\rightarrow$  p transition with transitions from the Xe inner 4d subshell, and how these CCR's change along the path Xe@C<sub>60</sub>  $\rightarrow$  Xe@C<sub>240</sub>  $\rightarrow$  Xe@C<sub>60</sub>@C<sub>240</sub>.

In performed calculations, fullerene cages were approximated by square-well potentials of certain inner radii  $r_0$ , widths  $\Delta$  and depths  $U_0$  [1, 3]. The Xe atom was placed at the center of fullerenes, both single-cage and nested fullerenes. The updated values of  $\Delta \approx 1.25$ ,  $r_0 \approx 6.01$  and  $U_0 \approx -0.422$  au [4] for C<sub>60</sub> were used in calculations. This is because they were shown [4] to provide a closer agreement between experiment

theory. As for C<sub>240</sub>,  $\Delta \approx 1.25$ ,  $r_0 \approx 12.875$  and  $U_0 \approx -0.52$  au [4]. Interchannel coupling was accounted for in the framework of the random phase approximation with exchange [5]. The calculated data are depicted in figure 1 where spectacular changes in the Xe 5s photoionization along the path Xe@C<sub>60</sub>  $\rightarrow$  Xe@C<sub>240</sub>  $\rightarrow$  Xe@C<sub>60</sub>@C<sub>240</sub> are seen. A detailed discussion will be offered at the spot.



**Figure 1.** Correlation confinement resonances in the Xe 5s photoionization cross section of Xe@C<sub>60</sub>, Xe@C<sub>240</sub> and Xe@C<sub>60</sub>@C<sub>240</sub>, as marked. They “mirror” CR's in the Xe 4d photoionization cross sections (not shown) of the endo-fullerenes in questions [4], due to interchannel coupling.

This work was supported by the NSF Grant No. PHY-0969386 and a UNA CoA&S grant.

## References

- [1] V. K. Dolmatov 2009 *Adv. Quant. Chem.* **58** 13
- [2] A. L. D. Kilcoyne *et al* 2010 *Phys. Rev. Lett.* **105** 213001
- [3] V. K. Dolmatov and S. T. Manson 2008 *J. Phys. B* **41** 165001
- [4] V. K. Dolmatov and D. A. Keating 2011 *XXVII ICPEAC* (Belfast, U.K., July 27 - Aug. 2, 2011) (submitted)
- [5] M. Ya. Amusia 1990 *Atomic Photoeffect* (Plenum Press, New York)

<sup>1</sup>E-mail: [vkdolmatov@una.edu](mailto:vkdolmatov@una.edu)